

Universality of the Distribution Functions of Random Matrix Theory

Craig A. Tracy

Department of Mathematics

and

Institute of Theoretical Dynamics

University of California, Davis, CA 95616, USA

e-mail address: tracy@itd.ucdavis.edu

Harold Widom

Department of Mathematics

University of California, Santa Cruz, CA 95064, USA

e-mail address: widom@math.ucsc.edu

February 9, 2008

Dedicated to James B. McGuire on the occasion of his sixty-fifth birthday.

1 Introduction

Statistical mechanical lattice models are called *solvable* if their associated Boltzmann weights satisfy the factorization or star-triangle equations of McGuire [1], Yang [2] and Baxter [3]. For such models the free energy per site and the one-point correlations in the thermodynamic limit are expressible in closed form [4]. There exists a deep mathematical structure [4, 5] underlying these solvable models; and near critical points, a wider applicability than one would initially expect. This last phenomenon, called *universality*, has its mathematical roots in the strong law of large numbers and the central limit theorems of probability theory and its physical origins in critical phenomena and conformal field theory.

The exact computation of n -point correlation functions is generally an open problem for most “solvable” models. In the special case of the 2D Ising model, the n -point functions (in the scaling limit) are expressible in terms of solutions to integrable differential equations [6, 7, 8]. The Wigner-Dyson theory of random matrices [9, 10] is a second class of statistical models where integrable differential equations and n -point correlations (or more precisely, level-spacing distributions) are related. This paper reviews some of these relationships.

In section 2 we define the basic objects of random matrix theory (RMT). In section 3 we recall the *bulk scaling limit* and the *edge scaling limit* and express various distribution functions in terms of Painlevé transcendents. In section 4 we discuss the universality of these distribution functions.

2 Random Matrix Models

In the Gaussian models [9, 10], the probability density that the eigenvalues lie in infinitesimal intervals about the points $x_1, \dots, x_N$ is given by

$$P_{N\beta}(x_1, \dots, x_N) = C_{N\beta} e^{-\frac{1}{2}\beta \sum x_i^2} \prod_{j < k} |x_j - x_k|^\beta, \quad (1)$$

where $C_{N\beta}$ is a normalization constant and

$$\beta := \begin{cases} 1 & \text{for GOE,} \\ 2 & \text{for GUE,} \\ 4 & \text{for GSE.} \end{cases}$$

We recall that for $\beta = 1$ the matrices are $N \times N$ real symmetric, for $\beta = 2$ the matrices are $N \times N$ complex Hermitian, and for $\beta = 4$ the matrices are $2N \times 2N$ self-dual Hermitian matrices. (For $\beta = 4$ each eigenvalue has multiplicity two.) In each case the eigenvalues are real.

In RMT the probabilities

$$\begin{aligned} E_{N\beta}(0; J) &:= \int_{x_j \notin J} \cdots \int P_{N\beta}(x_1, \dots, x_N) dx_1 \cdots dx_N, \\ &= \text{probability no eigenvalues lie in } J, \end{aligned}$$

are particularly interesting [10]. The simplest choices of J are (a, b) and (t, ∞) . In the first instance the mixed second partial derivative $E_{N\beta}(0; J)$ with respect to a and b gives the spacing distribution between consecutive eigenvalues; and in the second case, $F_{N\beta}(t) := E_{N\beta}(0; (t, \infty))$ is the distribution function for the largest eigenvalue.

The important fact [11, 12, 13] (see [14] for simplified proofs) is that $E_{N\beta}(0; J)$ (or its square for $\beta = 1$ or 4) is expressible as a Fredholm determinant. For $\beta = 2$ the kernel is a scalar kernel acting on J whereas for $\beta = 1, 4$, $E_{N\beta}(0; J)^2$ equals a Fredholm determinant of 2×2 matrix kernel acting on J .

3 Painlevé Representations

The representation of $E_{N\beta}(0; J)$, or its square, as a Fredholm determinant of an operator with kernel of a special form permits the determinant to be expressed in terms of Painlevé functions [15, 17, 18, 19].

3.1 Bulk Scaling Limit

Denote the density of eigenvalues at the point x_0 by $\rho(x_0)$. It is customary in the limit $N \rightarrow \infty$ to scale distances so that the resulting density is one. Precisely, we define $\xi = \rho_N(x_0)(x - x_0)$, x_0 independent of N , and consider the limit $N \rightarrow \infty$, $x \rightarrow x_0$, such that ξ is fixed. By requiring x_0 to be independent of N , we are choosing a point in the “bulk” of the spectrum and are examining the local statistics of the eigenvalues in some small neighborhood of the point x_0 . In this limit, and for $\beta = 2$, we are led to the Fredholm determinant of the operator on $L^2(0, s)$ whose kernel is the famous sine-kernel [11, 10]

$$K(\xi, \xi') := \frac{1}{\pi} \frac{\sin \pi(\xi - \xi')}{\xi - \xi'}.$$

Observe that the kernel is translationally invariant and independent of the point x_0 .

It is a result of Jimbo et al. [15] that

$$\det(I - \lambda K) = \exp \left(\int_0^{\pi s} \frac{\sigma(x; \lambda)}{x} dx \right),$$

where σ satisfies the differential equation

$$(x\sigma'')^2 + 4(x\sigma' - \sigma)(x\sigma' - \sigma + (\sigma')^2) = 0$$

with the boundary condition $\sigma(x; \lambda) \sim -\frac{\lambda}{\pi} x$ as $x \rightarrow 0$. (Here K is the operator whose kernel is the sine-kernel acting on $L^2(0, s)$.) The differential equation is the “ σ representation” of the P_V equation [15, 20]. For other proofs of this result see [21, 22, 16].

If $E_\beta(0; s)$ denotes the limiting value of $E_{N\beta}(0; (-t, t))$ in the bulk scaling limit with the scaled length of J set equal to s , then [10, 16, 19]

$$\begin{aligned} E_1(0; s) &= \det(I - K_+), \\ E_2(0; s) &= \det(I - K), \\ E_4(0; s/2) &= \frac{1}{2} (\det(I - K_+) + \det(I - K_-)) \end{aligned}$$

where $K_\pm$ are the operators with kernels $K(x, y) \pm K(-x, y)$. The $\det(I - K_\pm)$ can be expressed in terms of $\det(I - K)$ [15, 16]:

$$\det(I - K_\pm)^2 = \det(I - K) \exp \left(\mp \int_0^s \sqrt{-\frac{d^2}{dx^2} \log \det(I - K)} dx \right).$$

These formulas for $E_\beta(0; s)$ are well adapted for producing graphs of $E_\beta(0; s)$ and the level-spacing densities $p_\beta(s) = d^2 E_\beta(0; s)/ds^2$ once one knows the large s asymptotics of $\sigma(s; 1)$ and $E_\beta(0; s)$ [24, 16, 25].

3.2 Edge Scaling Limit

The limiting law, called the Wigner semi-circle law, is well known

$$\lim_{N_\beta \rightarrow \infty} \frac{1}{2\sigma\sqrt{N_\beta}} \rho_N(2\sigma\sqrt{N_\beta}x) = \begin{cases} \frac{1}{\pi} \sqrt{1-x^2} & |x| < 1, \\ 0 & |x| > 1. \end{cases}$$

Here σ , $\sigma/\sqrt{2}$, $\sigma/\sqrt{2}$ (for $\beta = 1, 2, 4$, respectively) is the standard deviation of the Gaussian distribution in the off-diagonal elements and

$$N_\beta = \begin{cases} N, & \beta = 1, \\ N, & \beta = 2, \\ 2N + 1, & \beta = 4. \end{cases}$$

For the normalization here, $\sigma = 1/\sqrt{2}$. Less known is that the distribution function of the largest eigenvalue satisfies [26]:

$$F_{N\beta}(2\sigma\sqrt{N_\beta} + x) \rightarrow \begin{cases} 0 & \text{if } x < 0, \\ 1 & \text{if } x > 0, \end{cases}$$

as $N \rightarrow \infty$. The edge scaling variable, s , defined by

$$t = 2\sigma\sqrt{N_\beta} + \frac{\sigma s}{N_\beta^{1/6}},$$

Table 1: The mean (μ_β), standard deviation (σ_β), skewness (S_β) and kurtosis (K_β) of F_β .

β	μ_β	σ_β	S_β	K_β
1	-1.20653	1.2680	0.293	0.165
2	-1.77109	0.9018	0.224	0.093
4	-2.30688	0.7195	0.166	0.050

gives the scale of the fluctuations at the edge of the spectrum [27, 28, 17]. As $N \rightarrow \infty$ with s fixed [17, 18]

$$F_{N2}(t) \rightarrow \exp \left(- \int_s^\infty (x-s)q(x)^2 dx \right) =: F_2(s) \quad (2)$$

where q is the solution to the P_{II} equation

$$q'' = s q + 2 q^3$$

satisfying the condition

$$q(s) \sim \text{Ai}(s) \quad \text{as } s \rightarrow \infty,$$

with Ai the Airy function.

For $\beta = 1, 4$ the results are [19]

$$F_{N1}(t)^2 \rightarrow F_1(s)^2 = F_2(s) \exp \left(- \int_s^\infty q(x) dx \right)$$

and


$$F_{N4}(t)^2 \rightarrow F_4(s/\sqrt{2})^2 = F_2(s) \cosh^2 \left(\frac{1}{2} \int_s^\infty q(x) dx \right).$$

Table 1 gives some statistics of F_β and Figure 1 shows the densities $f_\beta(s) = dF_\beta/ds$.

4 Universality

The universality of the distribution functions of RMT largely accounts for its success in a wide range of applications, see, e.g. [29]. However, this universality has only been *proved* in a small number of cases. The evidence for universality is largely numerical.

Figure 1: The probability density f_β of the largest eigenvalue, $\beta = 1, 2, 4$.



There are two types of universality results. The first, and the easier to establish rigorously, consists in modifying the random matrix model itself by, say, replacing the Gaussian $x^2/2$ appearing in (1) by an “arbitrary” potential $V(x)$. Changing the potential V changes the density of eigenvalues ρ , but several authors have established that the bulk scaling limit results in the sine-kernel and at soft edges one generically obtains the Airy universality class. (Fine tuning at the edge can result in different universality classes, see, e.g. [27].)

The second type of universality, and the one first envisioned by Wigner, asserts [30] that for a classical, “fully” chaotic Hamiltonian the corresponding quantum system has a level spacing distribution equal to $p_\beta(s)$ in the bulk. (The symmetry class determines which ensemble.) A nice numerical example of this *quantum chaos* is Robnik’s work [31] on chaotic billiards.

Here we discuss three other examples of RMT universality.


4.1 Zeros of the Riemann Zeta Function

Work by Montgomery [32] followed by extensive numerical calculations by Odlyzko [33] on zeros of the Riemann zeta function have given convincing numerical evidence that the normalized consecutive spacings follow the GUE distribution, see Figure 2 (the GUE Hypothesis). Rudnick and Sarnak [34] have proved a restricted form of this hypothesis.

4.2 Eigenvalues of Adjacency Matrices of Quasiperiodic Tilings

The discovery of quasicrystals has made the study of statistical mechanical models whose underlying lattice is quasiperiodic of considerable interest to

Figure 2: Data for nearest neighbor spacings among 1,041,600 zeros of the Riemann zeta function near the 2×10^{20} -th zero are plotted together with the GUE spacing density. Courtesy of Andrew Odlyzko [33].




physicists. In particular, in order to understand transport properties, tight binding models have been defined on various quasiperiodic lattices. One such study by Zhong et al. [35] defined a simplified tight binding model for the octagonal tiling of Ammann and Beenker. This quasiperiodic tiling consists of squares and rhombi with all edges of equal lengths (see, e.g., [35]) and has a D_8 symmetry around the central vertex. On this tiling the authors take as their Hamiltonian the adjacency matrix for the graph with free boundary conditions. The largest lattice they consider has 157,369 vertices. The matrix splits into ten blocks according to the irreducible representations of the dihedral group D_8 . For each of these ten independent subspectra, they compare the empirical cumulative distribution of the normalized spacings of the consecutive eigenvalues with the integrated density $I_1(s) = \int_s^\infty p_1(x) dx$ where p_1 is the GOE level spacing density. In Figure 3 we have reproduced a small portion of their data for one such subspectrum together with I_1 .

4.3 Distribution of the Length of the Longest Increasing Subsequence of Random Permutations

An old problem going back to Ulam asks for the limiting behavior of the length of the longest increasing subsequence of a random permutation. Precisely, if π is a permutation of $\{1, 2, \dots, N\}$, we say that $\pi(i_1), \dots, \pi(i_k)$ is an increasing

Figure 3: Data for nearest neighbor, cumulative, normalized spacings of eigenvalues of the adjacency matrix for a quasiperiodic octagonal tiling are plotted together with the GOE integrated spacing density $I_1(s)$. Data are from one independent subspectrum of a D_8 -symmetric octagonal patch of a tiling with 157,369 vertices. Courtesy of Zhong et al. [35].



subsequence in π if $i_1 < \dots < i_k$ and $\pi(i_1) < \dots < \pi(i_k)$. Let $\ell_N(\pi)$ be the length of the longest increasing subsequence of π . We take each permutation to be equally likely thus making ℓ_N a random variable. The problem then is to understand the distribution of ℓ_N . This problem has a long history (see, e.g. [36]) but its connection with RMT is recent [37, 38]. In a recent paper, Baik, Deift and Johansson [36] have proved the following remarkable result:

Theorem: Let S_N be the group of all permutations of N numbers with uniform distribution and let $\ell_N(\pi)$ be the length of the longest increasing subsequence of $\pi \in S_N$. Let χ be a random variable whose distribution function is F_2 , the distribution function for the largest eigenvalue in the GUE in the edge scaling limit (2). Then, as $N \rightarrow \infty$,

$$\chi_N := \frac{\ell_N - 2\sqrt{N}}{N^{1/6}} \rightarrow \chi$$

in distribution, i.e.

$$\lim_{N \rightarrow \infty} \text{Prob} \left(\chi_N := \frac{\ell_N - 2\sqrt{N}}{N^{1/6}} \leq s \right) = F_2(s)$$

for all $s \in \mathbf{R}$.

Acknowledgments

The authors wish to thank Andrew Odlyzko and Uwe Grimm for providing them with their data and allowing us to reproduce it here. We thank J. Baik for sending us [36] prior to publication. The first author acknowledges helpful conversations with Bruno Nachtergaele and Marko Robnik. This work was supported in part by the National Science Foundation through grants DMS-9802122 and DMS-9732687, and this support is gratefully acknowledged.

References

- [1] J. B. McGuire, *J. Math. Phys.* **5**, 622 (1964).
- [2] C. N. Yang, *Phys. Rev. Lett.* **19**, 1312 (1967).
- [3] R. J. Baxter, *Ann. Phys.* **70**, 193 (1972).
- [4] R. J. Baxter, *Exactly Solved Models in Statistical Mechanics* (Academic Press, NY, 1982).
- [5] M. Jimbo and T. Miwa, *Algebraic Analysis of Solvable Lattice Models*, CBMS Regional Conference Series in Mathematics, 85 (American Mathematical Society, Providence, 1995).
- [6] T. T. Wu, B. M. McCoy, C. A. Tracy and E. Barouch, *Phys. Rev.* **B13**, 316 (1976).
- [7] B. M. McCoy, C. A. Tracy and T. T. Wu, *J. Math. Phys.* **18**, 1058 (1977).
- [8] M. Sato, T. Miwa and M. Jimbo, *Publ. RIMS, Kyoto Univ.* **14**, 223 (1978); **15**, 201 (1979); **15**, 577 (1979); **15**, 871 (1979); **16**, 531 (1980).
- [9] C. E. Porter, *Statistical Theories of Spectra: Fluctuations* (Academic Press, NY, 1965).
- [10] M. L. Mehta, *Random Matrices*, 2nd ed. (Academic Press, San Diego, 1991).
- [11] M. Gaudin, *Nucl. Phys.* **25**, 447 (1961) (Reprinted in [9]).
- [12] F. J. Dyson, *Commun. Math. Phys.* **19**, 235 (1970).
- [13] M. L. Mehta, *Commun. Math. Phys.* **20**, 245 (1971).
- [14] C. A. Tracy and H. Widom, preprint. (solv-int/9804004).

- [15] M. Jimbo, T. Miwa, Y. Môri and M. Sato, *Physica D***1**, 80 (1980).
- [16] C. A. Tracy and H. Widom, in *Geometric and Quantum Aspects of Integrable Systems*, G. F. Helminck, Lect. Notes in Physics, Vol. 424, 103 (Springer-Verlag, Berlin, 1993).
- [17] C. A. Tracy and H. Widom, *Commun. Math. Phys.* **159**, 151 (1994).
- [18] C. A. Tracy and H. Widom, *Commun. Math. Phys.* **163**, 33 (1994).
- [19] C. A. Tracy and H. Widom, *Commun. Math. Phys.* **177**, 727 (1996).
- [20] M. Jimbo and T. Miwa, *Physica D***2**, 407 (1981).
- [21] M. L. Mehta, *J. de Phys. I France* **2**, 1721 (1992).
- [22] F. J. Dyson, in *Chen Ning Yang: A Great Physicist of the Twentieth Century*, eds. C. S. Liu and S.-T. Yau (International Press, Cambridge, 1995), pg. 131.
- [23] F. J. Dyson, *Commun. Math. Phys.* **47**, 171 (1976).
- [24] E. L. Basor, C. A. Tracy and H. Widom, *Phys. Rev. Lett.* **69**, 5 (1992).
- [25] F. V. Andreev and A. V. Kitaev, *Math. Res. Lett.* **4**, 741 (1997).
- [26] Z. D. Bai and Y. Q. Yin, *Ann. Prob.* **16**, 1729 (1988).
- [27] M. J. Bowick and E. Brézin, *Phys. Letts. B***268**, 21 (1991).
- [28] P. J. Forrester, *Nucl. Phys. B***402**[FS], 709 (1993).
- [29] T. Guhr, A. Mueller-Groeling, H. A. Weidenmueller, *Phys. Rept.* **299**, 189 (1998).
- [30] O. Bohigas, M. J. Giannoni and C. Schmit, *Phys. Rev. Lett.* **52**, 1 (1984).
- [31] M. Robnik, *J. Phys. A***25**, 3593 (1992).
- [32] H. L. Montgomery, in *Analytic Number Theory*, Proc. Sympos. Pure Math **24**, 181 (1973).
- [33] A. M. Odlyzko, *Math. Comp.* **48**, 273 (1987).
- [34] Z. Rudnick and P. Sarnak, *Duke Math. J.* **81**, 269 (1996).

- [35] J. X. Zhong, U. Grimm, R. A. Römer and M. Schreiber, *Phys. Rev. Lett.* **80**, 3996 (1998). [See also M. Schreiber, U. Grimm, R. A. Römer and J. X. Zhong, Application of random matrix theory to quasiperiodic systems, to appear in the proceedings of Percolation98.]
- [36] J. Baik, P. Deift, K. Johansson, preprint.
- [37] A. M. Odlyzko, B. Poonen, H. Widom and H. S. Wilf, unpublished manuscript.
- [38] E. M. Rains, *Elect. J. Combinatorics* **5**, R12 (1998).